

Non-commutative Hardy inequalities

Frank Hansen

June 4, 2008

Revised version January 8, 2009

Abstract

We extend Hardy's inequality from sequences of non-negative numbers to sequences of positive semi-definite operators if the parameter p satisfies $1 < p \leq 2$, and to operators under a trace for arbitrary $p > 1$. Applications to trace functions are given. We introduce the tracial geometric mean and generalize Carleman's inequality.

Key words and phrases: Hardy's inequality, positive operator, trace function, geometric mean, Carleman's inequality.

1 Introduction

The standard form of Hardy's inequality [7, 8, 9, 10] asserts that

$$(1) \quad \sum_{n=1}^{\infty} \left(\frac{1}{n} \sum_{k=1}^n a_k \right)^p \leq \left(\frac{p}{p-1} \right)^p \sum_{n=1}^{\infty} a_n^p$$

for $p > 1$ and any sequence $a = (a_n)$ of non-negative real numbers in l_p . The inequality is sharp in the sense that the constant $(p/(p-1))^p$ cannot be replaced by a smaller number such that the inequality remains true for all (even finite) sequences of non-negative real numbers.

The arithmetic averaging operator h (or the Hardy operator) is defined by setting

$$(2) \quad h(a) = \left(\frac{1}{n} \sum_{k=1}^n a_k \right)_{n=1}^{\infty}$$

for any sequence $a = (a_1, a_2, \dots)$ of real or complex numbers. Hardy's inequality thus states that h maps l_p into itself and has norm $p/(p-1)$.

We shall prove (1) also for sequences of positive semi-definite operators on a Hilbert space provided $1 < p \leq 2$. Under the trace the inequality is valid for arbitrary $p > 1$ and (appropriate) sequences of positive semi-definite operators. The classical proofs of Hardy's inequality are not easy to extend to operators. The key to the results in this paper is the realization that, as is the case for many other classical inequalities, Hardy's inequality is in essence an expression of convexity. The author is indebted to Lars-Erik Persson who in a talk mentioned that there is a connection between convexity and Hardy's inequality, and that it is part of the unpublished folklore known to specialist in classical inequalities.

2 Non-commutative generalizations

Lemma 2.1. *Let $1 \leq p \leq 2$ be a real number, and let $g: (0, \infty) \rightarrow B(H)_+$ be a weakly measurable map such that the integral*

$$\int_0^\infty g(x)^p \frac{dx}{x}$$

defines a bounded linear operator on H . Then the operator inequality

$$\int_0^\infty \left(\frac{1}{x} \int_0^x g(t) dt \right)^p \frac{dx}{x} \leq \int_0^\infty g(x)^p \frac{dx}{x}$$

is valid. The inequality cannot (generally) be improved by inserting a constant less than one in front of the second integral.

Proof. Since the function $t \rightarrow t^p$ is operator convex we obtain

$$\left(\frac{1}{x} \int_0^x g(t) dt \right)^p \leq \frac{1}{x} \int_0^x g(t)^p dt \quad x > 0$$

with equality for functions g which are constant in $(0, x]$. Consequently

$$\begin{aligned} \int_0^\infty \left(\frac{1}{x} \int_0^x g(t) dt \right)^p \frac{dx}{x} &\leq \int_0^\infty \frac{1}{x} \int_0^x g(t)^p dt \frac{dx}{x} \\ &= \int_0^\infty g(t)^p \int_t^\infty \frac{1}{x^2} dx dt = \int_0^\infty g(t)^p \frac{dt}{t} \end{aligned}$$

and the assertions are proved. **QED**

Lemma 2.2. *Let $p \geq 1$ be a real number, and let $g: (0, \infty) \rightarrow B(H)_+$ be a weakly measurable map such that the integral*

$$\int_0^\infty g(x)^p \frac{dx}{x}$$

defines a bounded linear trace class operator on H . Then the inequality

$$\mathrm{Tr} \int_0^\infty \left(\frac{1}{x} \int_0^x g(t) dt \right)^p \frac{dx}{x} \leq \mathrm{Tr} \int_0^\infty g(x)^p \frac{dx}{x}$$

is valid. The inequality cannot (generally) be improved by inserting a constant less than one in front of the second integral.

Proof. The function $t \rightarrow t^p$ is for $p \geq 1$ convex in operators under a trace. That is, the function $X \rightarrow \mathrm{Tr} X^p$ is convex in positive semi-definite operators X such that X^p is trace class. The same steps as in the proof of the preceding lemma, mutatis mutandis, may be carried out and the assertions follow. **QED**

Theorem 2.3. *Let $1 < p \leq 2$ be a real number, and let $f: (0, \infty) \rightarrow B(H)_+$ be a weakly measurable map such that the integral*

$$\int_0^\infty f(x)^p dx$$

defines a bounded linear operator on H . Then the inequality

$$\int_0^\infty \left(\frac{1}{x} \int_0^x f(t) dt \right)^p dx \leq \left(\frac{p}{p-1} \right)^p \int_0^\infty f(x)^p dx$$

holds, and the constant $(p/(p-1))^p$ is the best possible.

Proof. We apply the function

$$g(t) = f(t^{p/(p-1)}) t^{1/(p-1)}$$

in Lemma 2.1 and obtain

$$\int_0^\infty \left(\frac{1}{x} \int_0^x f(t^{p/(p-1)}) t^{1/(p-1)} dt \right)^p \frac{dx}{x} \leq \int_0^\infty f(x^{p/(p-1)})^p x^{p/(p-1)} \frac{dx}{x}$$

Setting $y = t^{p/(p-1)}$ and thus

$$\frac{dy}{dt} = \frac{p}{p-1} \cdot t^{1/(p-1)}$$

in the first integral, we obtain

$$\begin{aligned} & \int_0^\infty \left(\frac{1}{x} \int_0^x f(t^{p/(p-1)}) t^{1/(p-1)} dt \right)^p \frac{dx}{x} \\ &= \left(\frac{p-1}{p} \right)^p \int_0^\infty \left(\frac{1}{x} \int_0^{x^{p/(p-1)}} f(y) dy \right)^p \frac{dx}{x} \end{aligned}$$

and thus

$$\left(\frac{p-1}{p} \right)^p \int_0^\infty \left(\frac{1}{x} \int_0^{x^{p/(p-1)}} f(y) dy \right)^p \frac{dx}{x} \leq \int_0^\infty f(x^{p/(p-1)})^p x^{p/(p-1)} \frac{dx}{x}.$$

We finally set $z = x^{p/(p-1)}$ and thus

$$\frac{dz}{dx} = \frac{p}{p-1} \cdot \frac{z}{x}$$

in both integrals and obtain

$$\left(\frac{p-1}{p} \right)^{p+1} \int_0^\infty \left(\frac{1}{z^{(p-1)/p}} \int_0^z f(y) dy \right)^p \frac{dz}{z} \leq \frac{p-1}{p} \int_0^\infty f(z)^p dz.$$

This is rearranged to

$$\int_0^\infty \left(\frac{1}{z} \int_0^z f(y) dy \right)^p dz \leq \left(\frac{p}{p-1} \right)^p \int_0^\infty f(z)^p dz$$

and the assertions are proved. **QED**

Theorem 2.4. *Let $1 < p$ be a real number, and let $f: (0, \infty) \rightarrow B(H)_+$ be a weakly measurable map such that the integral*

$$\int_0^\infty f(x)^p dx$$

defines a bounded linear trace class operator on H . Then the inequality

$$\text{Tr} \int_0^\infty \left(\frac{1}{x} \int_0^x f(t) dt \right)^p dx \leq \left(\frac{p}{p-1} \right)^p \text{Tr} \int_0^\infty f(x)^p dx$$

holds, and the constant $(p/(p-1))^p$ is the best possible.

Proof. We apply for $p > 1$ the function

$$g(t) = f(t^{p/(p-1)})t^{1/(p-1)}$$

in Lemma 2.2 and make the same substitutions under the trace as applied in the proof of Theorem 2.3. **QED**

By considering the case where f is a step function we obtain:

Corollary 2.5. *Let $1 < p \leq 2$ be a real number, and let $a = (a_1, a_2, \dots)$ be a sequence of positive semi-definite operators on a Hilbert space such that the infinite sum $a_1^p + a_2^p + \dots$ defines a bounded operator. Then the inequality*

$$(3) \quad \sum_{n=1}^{\infty} \left(\frac{1}{n} \sum_{k=1}^n a_k \right)^p \leq \left(\frac{p}{p-1} \right)^p \sum_{n=1}^{\infty} a_n^p$$

holds, and the constant $(p/(p-1))^p$ is the best possible.

Corollary 2.6. *Let $1 < p$ be a real number, and let $a = (a_1, a_2, \dots)$ be a sequence of positive semi-definite operators on a Hilbert space such that the infinite sum $a_1^p + a_2^p + \dots$ defines a bounded trace class operator. Then the inequality*

$$(4) \quad \text{Tr} \sum_{n=1}^{\infty} \left(\frac{1}{n} \sum_{k=1}^n a_k \right)^p \leq \left(\frac{p}{p-1} \right)^p \text{Tr} \sum_{n=1}^{\infty} a_n^p$$

holds, and the constant $(p/(p-1))^p$ is the best possible.

2.1 Trace functions

Consider for $p > 0$ the homogeneous functional

$$\Phi_p(a) = \text{Tr} \left[\left(\sum_{n=1}^{\infty} a_n^p \right)^{1/p} \right]$$

defined on sequences $a = (a_1, a_2, \dots)$ of positive definite operators acting on a Hilbert space such that $(a_1^p + a_2^p + \dots)^{1/p}$ defines a bounded trace class operator. It is a consequence of Epstein's theorem that Φ_p is concave, cf. [5] and [4, Theorem 1.1]. Carlen and Lieb conjectured in 1999 that Φ_p is

convex for $1 \leq p \leq 2$, and they showed that this would lead to a Minkowski inequality for operators on a tensor product of three Hilbert spaces and to another proof of strong subadditivity of entropy. Carlen and Lieb eventually proved the conjecture in 2008 [4, Theorem 1.1].

Since Φ_p is convex for $1 \leq p \leq 2$ the domain is a convex cone, but the functional cannot be extended to a norm. The obstacle is that the modulus $|X|$ of an operator X is not a convex function in X .

Theorem 2.7. *Let $1 < p \leq 2$ be a real number, and let $a = (a_1, a_2, \dots)$ be a sequence of positive semi-definite operators on a Hilbert space such that $(a_1^p + a_2^p + \dots)^{1/p}$ defines a bounded trace class operator. Then*

$$\Phi_p(h(a)) \leq \frac{p}{p-1} \Phi_p(a),$$

where h is the Hardy operator (2).

Proof. Since $t \rightarrow t^\alpha$ is operator monotone for $0 \leq \alpha \leq 1$, it follows from the non-commutative Hardy inequality (3) that

$$\left(\sum_{n=1}^{\infty} \left[\frac{1}{n} \sum_{k=1}^n a_k \right]^p \right)^{1/p} \leq \frac{p}{p-1} \left(\sum_{n=1}^{\infty} a_n^p \right)^{1/p}$$

for $1 < p \leq 2$. The assertion then follows by taking the trace. **QED**

3 The tracial geometric mean

The operator power mean¹ is defined by

$$M_p(a_1, \dots, a_n) = \left(\frac{1}{n} \sum_{k=1}^n a_k^{1/p} \right)^p.$$

Let now $q \geq p \geq 1$. Since $t \rightarrow t^{p/q}$ is operator concave we obtain

$$M_p(a_1, \dots, a_n)^{1/q} = \left(\frac{1}{n} \sum_{k=1}^n a_k^{1/p} \right)^{p/q} \geq \frac{1}{n} \sum_{k=1}^n a_k^{1/q} = M_q(a_1, \dots, a_n)^{1/q},$$

¹Sometimes p is replaced with $1/p$ in the literature [6].

and since $t \rightarrow t^q$ is monotone under a trace, that is $X \rightarrow \text{Tr } X^q$ is monotone increasing in positive definite X , we also obtain

$$\text{Tr } M_p(a_1, \dots, a_n) \geq \text{Tr } M_q(a_1, \dots, a_n).$$

The trace of the operator mean is thus a decreasing function in the power parameter p , so the following definition makes sense:

Definition 3.1. *We define the tracial geometric mean by setting*

$$TG(a_1, \dots, a_n) = \lim_{p \rightarrow \infty} \text{Tr} \left(\frac{1}{n} \sum_{k=1}^n a_k^{1/p} \right)^p$$

for positive semi-definite trace class operators $a_1, \dots, a_n$ on a Hilbert space.

Notice (by setting $p = 1$) the inequality

$$\text{Tr} \left(\frac{a_1 + \dots + a_n}{n} \right) \geq TG(a_1, \dots, a_n).$$

The tracial geometric mean has a number of attractive properties:

Theorem 3.2. *Let $a_1, \dots, a_n$ be positive semi-definite bounded operators on a Hilbert space and consider the tracial geometric mean $TG(a_1, \dots, a_n)$.*

- (1) *It coincides with the geometric mean $(a_1 \cdots a_n)^{1/n}$ for positive numbers and with $\text{Tr} (a_1 \cdots a_n)^{1/n}$ for commuting operators.*
- (2) *It is a (tracial) mean; $TG(a, \dots, a) = \text{Tr } a$.*
- (3) *It is homogeneous; $TG(ta_1, \dots, ta_n) = t \cdot TG(a_1, \dots, a_n)$ for $t > 0$.*
- (4) *It is symmetric in the entries.*
- (5) *It is unitarily invariant.*
- (6) *It acts additively on sequences of block matrices.*
- (7) *It is (separately) increasing in the entries.*
- (8) *It is (jointly) concave in the entries.*

Proof. The first statement for positive numbers is a classical result and (2) to (6) are quite obvious.

To prove (7) we notice that $M_p(a_1, \dots, a_n)^{1/p}$ is separately increasing in the entries by operator monotonicity of $t \rightarrow t^{1/p}$. We therefore obtain, as above, that $\text{Tr } M_p(a_1, \dots, a_n)$, for each $p \geq 1$, is separately increasing in the entries. This property is hence satisfied also for the tracial geometric mean by taking the limit $p \rightarrow \infty$.

To prove (8) we notice that $\text{Tr } M_p(a_1, \dots, a_n)$, for each $p \geq 1$, is a concave function in the entries by Epstein's theorem, cf. [5] and [4, Theorem 1.1]. Taking the limit $p \rightarrow \infty$ we therefore realize that also the tracial geometric mean is a concave function in the entries. **QED**

The last property (8) is by far the most difficult to establish. The concrete formula given in the next theorem is related to [6, Theorem 4.19]. It is interesting to notice that it is of no use to prove (8) in the previous result.

Theorem 3.3. *If the operators $a_1, \dots, a_n$ are strictly positive, then the tracial geometric mean*

$$(5) \quad TG(a_1, \dots, a_n) = \text{Tr} \exp \left(\frac{1}{n} \sum_{k=1}^n \log a_k \right).$$

Proof. Since the logarithm is operator concave, we have

$$p \cdot \log \left(\frac{1}{n} \sum_{k=1}^n a_k^{1/p} \right) \geq \frac{1}{n} \sum_{k=1}^n \log a_k$$

and hence under the trace

$$\begin{aligned} M_p(a_1, \dots, a_n) &= \text{Tr} \exp \left[p \cdot \log \left(\frac{1}{n} \sum_{k=1}^n a_k^{1/p} \right) \right] \\ &\geq \text{Tr} \exp \left(\frac{1}{n} \sum_{k=1}^n \log a_k \right) \end{aligned}$$

for any $p \geq 1$. Taking the limit $p \rightarrow \infty$ we thus obtain that the left hand side in (5) majorizes the right hand side. To verify the opposite inequality

we use $\log t \leq t - 1$ to obtain

$$\begin{aligned} \log M_p(a_1, \dots, a_k) &= p \cdot \log \left(\frac{1}{n} \sum_{k=1}^n a_k^{1/p} \right) \\ &\leq \frac{1}{n} \sum_{k=1}^n p(a_k^{1/p} - 1) \end{aligned}$$

and hence under the trace

$$TG(a_1, \dots, a_n) \leq \text{Tr } M_p(a_1, \dots, a_n) \leq \text{Tr} \exp \left(\frac{1}{n} \sum_{k=1}^n p(a_k^{1/p} - 1) \right)$$

for any $p \geq 1$. Since $p(a_k^{1/p} - 1) \rightarrow \log a_k$ in the strong operator topology for $p \rightarrow \infty$, the assertion follows. **QED**

The following theorem is a non-commutative version of Carleman's inequality [3].

Theorem 3.4. *The inequality*

$$\sum_{n=1}^{\infty} TG(a_1, \dots, a_n) \leq e \sum_{n=1}^{\infty} \text{Tr } a_n$$

is valid for sequences of positive semi-definite trace class operators such that the sum on the right hand side is finite.

Proof. If we in the tracial Hardy inequality (4) replace a_n with $a_n^{1/p}$ for each n then we obtain

$$\text{Tr} \sum_{n=1}^{\infty} \left(\frac{1}{n} \sum_{k=1}^n a_k^{1/p} \right)^p \leq \left(\frac{p}{p-1} \right)^p \text{Tr} \sum_{n=1}^{\infty} a_n$$

for $p > 1$. Since

$$\left(\frac{p}{p-1} \right)^p \rightarrow e \quad \text{for } p \rightarrow \infty$$

the statement follows by letting $p \rightarrow \infty$ in the inequality. **QED**

References

- [1] T. Ando, C.-K. Li, and R. Mathias. Geometric means. *Linear Algebra Appl.*, 385:305–334, 2004.
- [2] R. Bhatia and J. Holbrook. Noncommutative geometric means. *Math. Intelligencer*, 28:32–39, 2006.
- [3] T. Carleman. Sur les fonctions quasi-analytiques. Fifth Scandinavian Congress of Mathematicians (Helsinki, 1922), pages 181–197. Akadem. Buchh., 1923.
- [4] E.A. Carlen and E.H. Lieb. A Minkowsky type trace inequality and strong subadditivity of quantum entropy II: Convexity and concavity. *Lett. Math. Phys.*, 83:107–126, 2008.
- [5] H. Epstein. Remarks on two theorems of E. Lieb. *Comm. Math. Phys.*, 31:317–325, 1973.
- [6] T. Furuta, Hot J.M., J. Pečarić, and Y. Seo. *Mond-Pečarić Method in Operator Inequalities*. Element, Zagreb, 2005.
- [7] G.H. Hardy. Note on a theorem of Hilbert. *Math. Z.*, 6:314–317, 1920.
- [8] G.H. Hardy. Notes on some points in the integral calculus, LX. An inequality between integrals. *Messenger of Math.*, 54:150–156, 1925.
- [9] G.H. Hardy, J.E. Littlewood, and G. Pólya. *Inequalities, 2nd ed.* Cambridge University Press, Cambridge, 1967.
- [10] A. Kufner, L. Maligranda, and L.-E. Persson. The prehistory of the Hardy inequality. *American Math. Month.*, pages 715–732, October 2006.
- [11] A. Kufner and L.-E. Persson. *Weighted inequalities of Hardy type*. World Scientific, Singapore, 2003.

Frank Hansen: Department of Economics, University of Copenhagen, Studiestraede 6, DK-1455 Copenhagen K, Denmark.